

INTERNATIONAL SOCIETY FOR CRANIOFACIAL SURGERY

X th International Congress
Monterey, California



Management of Upper Airway and
Feeding Difficulties in Infants with
Pierre Robin Sequence Using
Mandibular Distraction
Osteogenesis: Presenting a New
Protocol

**Children's Hospital of Michigan
Detroit Medical Center/Wayne State University**

Divisions of Plastic & Reconstructive Surgery

Joseph A. Broujerdi, M.D., D.M.D.

Arlene A. Rozzelle, M.D.



**Children's Hospital
of Michigan**

Detroit Medical Center/Wayne State University

Infant Mandibular Distraction

- 13 infants with PRS with or without cleft palate or other anomalies, 7 males & 6 females
 - 9 Cleft palate
 - 1 Cleft lip
 - 2 Treacher Collins Synd.
 - 1 Stickler Synd.
 - 1 Trisomy 18
- Grade II-III Caouette-Laberge airway-feeding classification (3II, 10 III)
 - I Prone & no feeding difficulty
 - II Prone & NG feeding only
 - III ETT with NGT (FTT),
(2 Tracheostomy, 1 G-tube)
- Age at distraction 8 do - 16 mo (average 4 mo)

Infant Mandibular Distraction

- Preoperative workup
 - 3D CT
 - Sleep Study
 - GER Scan
 - » 9 infants dx with GERD
 - DLB
 - » 4 infants dx with Laryngomalacia
 - » 1 infant dx with Tracheomalacia
 - Genetics consultation
 - ECHO Cardiogram
 - » 5 infants with congenital heart defect

Surgical Technique

- General anesthesia
- Submandibular incisions
- Corticotomies
 - » At the angle of the mandible
 - » Avoid IAN and tooth buds
- Insertion of Neonatal Distractor with micro monocortical screws at the inferior border of mandible
 - » Neonatal KLS-Martin
 - » Logic Jr. OsteoMed
- Wound closed in layers
- Distraction in AP vector



International Partners

Post Op Care

- Admission to ICU
- Distraction completed during hospital course
- Progress followed with mandibular x-rays
- Ultrasound for assessment of consolidation
- Post-Op sleep study
- Post-Op 3D Craniofacial CT after removal of distractors


Distraction Technique


- Latency period of 0-2 days
- Distraction rate of 1.0mm-1.5mm a day
- Distraction to a correct anatomical skeletal class I or class III in severe cases of OSA
- Distraction length 8mm-15mm (11mm)
- Consolidation period 8 wks-20 wks (10 wks)




Complication



- One marginal mandibular nerve paresis
- One readmission for pneumonia
- One distractor backed up











AVERAGE MANDIBULAR MEASUREMENT

■ Pre-distraktion

- Co-Go 17 mm
- Go-Pg 23 mm

■ Post-distraktion

- Co-Go 20 mm
- Go-Pg 32 mm

Vertical + 3 mm

Horizontal + 9 mm

SLEEP STUDY DATA

	Pre-Op AHI	Post-Op AHI
BG	47	11
CN	9	4
GB	30	9
NJ	37	10
PI	13	2
RR	18	5
SD	37	5
WK	44	12

SLEEP STUDY DATA

- Pre-op average AHI and lowest saturation

- » AHI 28
- » Sat. 73%

- Post-op average AHI and lowest saturation

- » AHI 8
- » Sat. 85%

Feeding & Weight Pattern Post Operative

■ Feeding

- 10 infants with good PO intake
- 2 infants with poor to moderate PO intake during the day & NG feeding during sleep
- 1 infant poor PO intake with G-tube feeding

■ Weight gain

- 11 good
- 1 moderate
- 1 poor

Sleep Pattern Post Operative

- 2 tracheostomies decanulated
- 11 patients sleep tolerating room air with no complaints from parents
- 1 patient sleeps on room air with pulse oximeter with infrequent desaturations
- 1 patient sleeps on O₂ via nasal cannula and pulse oximeter with infrequent desaturations

Conclusion

- Infant mandibular distraction is a feasible, effective option
 - » Low risk
 - » Technically easy
 - » Well tolerated
 - » Very effective