
IMPLANT PLACEMENT POST-OPERATIVE INSTRUCTIONS

1. **Pain control recommendations:** Your first level of pain control will be ibuprofen. You will be given 600mg of ibuprofen at the beginning of your appointment. It is recommended that you continue to take the ibuprofen for the next 3-4 days then as needed (600mg every 6 hours). The maximum dose of ibuprofen is 2400mg per day. Your second level of pain control is acetaminophen "Tylenol". Two 500mg tablets can be taken 3 hours after the initial dose of ibuprofen and continue to alternate these medications every 3 hours. Most patients are comfortable following these guidelines; however, should you need stronger pain relief you will be given an additional prescription to fill at the pharmacy. All of the prescription pain relievers contain acetaminophen: therefore, if you take the pain prescription medication, do not take any additional acetaminophen for 6 hours. **IMPORTANT:** The prescription pain medication may cause drowsiness or nausea. Do not drive or drink alcoholic beverages while taking the prescription pain medication. To avoid nausea, take with food or milk.
2. **Peridex mouth wash:** Apply to surgery area only. Do not rinse with Peridex. Apply by dipping soft toothbrush (provided) or q-tip into small cup and lightly brush around teeth and onto packing. Peridex may cause a temporary stain; To avoid as much stain as possible; Brush and floss non-surgery areas before using Peridex. Avoid smoking and excess amounts of coffee and or tea.
3. **Antibiotic prescription:** You will be given a prescription for an antibiotic at the time of your appointment. It is important that you take this prescription to completion.
4. The anesthetic used will typically last 4-6 hours. Be careful not to bite your tongue while you are still numb. Avoid hot food or drinks that will burn numb areas.
5. Soft foods should be eaten the first day or two. Avoid popcorn or foods with husks or shells for several weeks.
6. **Important:** Avoid any type of chewing, biting, or pressure on an implant for the entire time of healing (4-6 months). An implant is most vulnerable the first 3-4 weeks after placement. Therefore, extra caution should be used during this time.
7. A small amount of bleeding is considered normal the day of and the morning after surgery. Avoid rinsing your mouth vigorously or spitting as this will prolong bleeding.
8. If you are not already on a calcium supplement, a dose of 1000mg to 1500 mg of calcium per day for the duration of healing (4-6 months) is recommended.
9. **Ice is very helpful in reducing swelling:** Apply an ice pack to the surgery area (wrapped in a towel on the outside of the face) for 20 minutes at a time every hour the day of surgery.
10. **Avoid smoking**

Special Instructions

Should you have any questions or concerns call 344-0908

To reach Dr. Ferguson after hours call 484-2235.

To reach Dr. Beaudry after hours call (480) 444-8119.

Protect more than your Dental Health • ***Protect your Total Health!***